OCEAN MINERALS FORWARD »

Responsible Development: What the Science Shows

More Than 50 Years of Research Provides a Strong Foundation

Responsible ocean mineral development starts with understanding the environment. More than five decades of deep-ocean research — from NOAA's pioneering studies in the 1970s to today's highly instrumented pilot trials — have built an increasingly detailed scientific foundation for understanding and managing ocean mineral development.

That science allows developers and regulators to **establish environmental baselines before operations begin, monitor activity as it occurs, measure actual performance, and adapt technology and operations as knowledge advances.**

50+ YEARS
of deep-ocean environmental research

9,000+ DAYS
of research at sea

HUNDREDS
of deep-ocean research expeditions

Science Has Been Part of the Industry from the Beginning

Ocean mineral technology and environmental research have developed side by side.

1975–1980

NOAA DOMES

NOAA's Deep Ocean Mining Environmental Study established an early scientific framework for evaluating deep-seabed mineral development and environmental conditions.

1976–1979

Pioneer Collection Trials

Four U.S.-based consortia conducted multiple in-field nodule collection tests at depths of several thousand meters in the Clarion-Clipperton Zone, generating real-world engineering and environmental data.

2021

Patania II Trial

Independent scientists deployed roughly 200 instruments across 43 platforms during DEME's collector trial — monitoring the operation with an extensive network of sensors and scientific equipment.

2022

Integrated Pilot Collection

During The Metals Company's pilot collection trial, independent researchers monitored an integrated deep-sea collection system in real time, adding another major body of empirical data to the scientific record.

TODAY

Better Technology, Better Data

Autonomous vehicles, high-resolution mapping, advanced sensors, cameras, machine vision, and real-time monitoring are dramatically expanding what scientists can observe and measure.

What Real-World Research Is Showing

Field research allows scientists to test predictions against actual operations. Studies of modern collection trials have found that some measured effects differ significantly from earlier assumptions and modeling. Research associated with MIT, for example, found sediment disturbance from conventional collection was **one-third to one-sixth of some earlier estimates**, while observed sediment plumes remained much closer to the seafloor than some earlier predictions anticipated.

Long-term studies are equally important. Scientists have returned to historic disturbance sites decades later, documenting **recovery of key biological communities** while continuing to study changes that persist over longer periods. The scientific record continues to grow with every baseline survey, technology test, pilot program, and long-term study.

Sediment Plumes

Sediment movement is one of the most studied aspects of polymetallic nodule collection. Modern field trials are allowing researchers to replace assumptions with direct measurements.

Measurements also found that much of the sediment mobilized by the collector remained close to the seafloor rather than rising hundreds of feet through the water column as some earlier scenarios had anticipated.

Fisheries

Many polymetallic nodule resources occur 4,000–5,500 meters below the ocean surface, while major commercial fisheries operate much higher in the water column. That substantial vertical separation is an important feature of nodule development.

Environmental baseline studies and water-column monitoring can characterize species distribution, currents and other conditions before operations begin. During operations, monitoring can track conditions throughout the water column, while vessel coordination and project design can further reduce interactions with fishing activity.

Seafloor Disturbance


Polymetallic nodules lie unattached on the surface of the abyssal plain. Recovering them does not require drilling into rock, blasting an ore body, or removing conventional mine overburden. Modern collection systems are designed around controlled operations within defined collection areas, and newer autonomous and selective technologies offer additional opportunities to reduce sediment disturbance and seafloor contact.

Long-term studies of historic disturbance sites also allow researchers to observe what happens over decades. Those studies have documented recovery of key biological communities while continuing to identify and study effects that persist longer.

Biodiversity

Understanding biodiversity begins before collection. High-resolution imagery, genetic analysis, physical sampling, autonomous vehicles and other scientific tools can document species, habitats and biological communities across potential operating areas.

That creates something essential: a **before-and-after record.**



AMERICAN RESOURCES. AMERICAN INNOVATION. AMERICAN INDUSTRY.

Explore the full report and supporting resources at OceanMineralsForward.com

SCAN TO SEE FULL REPORT ONLINE